

Quantum Correction of Fluctuation Theorem

T. Monnai , S. Tasaki

*Department of Applied Physics , Waseda University, 3-4-1 Okubo,
Shinjuku-ku, Tokyo 169-8555, Japan*

Abstract

Quantum analogues of the transient and steady-state fluctuation theorems are investigated for a harmonic oscillator linearly coupled with a harmonic reservoir. The probability distribution for the work done externally is derived and the following facts are shown: (i) In the transient fluctuation theorem, there appears a quantum correction of order $\hbar^2$. (ii) In the steady-state fluctuation theorem, the existence of a quantum correction depends on the way of driving. In the uniformly dragged case, the classical formula holds, while, in the periodically driven case, there appears a correction of order $\hbar^2$.

Key words: Fluctuation theorem, Nonequilibrium fluctuation, Quantum corrections

PACS: 05.30.-d, 05.40.Ca

1 INTRODUCTION

In the linear nonequilibrium regime, statistical properties of fluctuations are known to characterize nonequilibrium states via exact identities such as the Green-Kubo formula and Fluctuation-Dissipation Theorems. On the contrary, general properties of the far-from-equilibrium fluctuations have not been well understood. Recently, two new identities valid even far from equilibrium were found: the fluctuation theorem (FT) first found by Evans, Cohen and Morriss[1] and the nonequilibrium free-energy equality given by Jarzynski[2].

The fluctuation theorem addresses the symmetry of entropy production or work externally done and several different versions are known. Nevertheless,

Email address: monnai@suou.waseda.jp (T. Monnai).

it takes the following general form:

$$\left(\lim_{\tau \rightarrow +\infty} \right) \frac{1}{\tau} \log \frac{\text{Prob}(\sigma)}{\text{Prob}(-\sigma)} = \sigma , \quad (1)$$

where $\text{Prob}(\sigma)$ stands for the probability of observing an average entropy production rate (or an average power applied externally), σ , during the time interval τ . The $\tau \rightarrow +\infty$ limit may or may not be taken depending on the situations.

FT with $\tau \rightarrow +\infty$ limit was first found numerically by Evans, Cohen and Morriss[1] for nonequilibrium steady states of thermostatted systems. Then, it was rigorously proved by Gallavotti and Cohen[3]. This version is referred to as the steady-state fluctuation theorem (SSFT). On the other hand, before the proof by Gallavotti and Cohen, Evans and Searles[4] derived the relation (1) without $\tau \rightarrow +\infty$ limit for transient trajectories starting from initial states obeying the microcanonical distribution. This version is referred to as the transient fluctuation theorem (TFT) and has been extensively studied by Evans et al.[5]. Moreover, it was extended to stochastically driven systems[6,7,8] and to open conservative systems[9,10,11]. Its relation with Jarzynski equality was studied as well[12] and the related topics have been extensively investigated (see e.g., references in Refs.[11,13]).

Recently, a beautiful experiment by Wang et al.[14] confirmed TFT for a colloidal particle kept in a uniformly moving optical trap. The dynamics of the colloidal particle is governed by the Langevin equation and TFT holds for the work done externally. FT for the Langevin equation was studied by Mazonka and Jarzynski[15] to illustrate the difference between SSFT and TFT, and the experiment stimulates the reinvestigations of FT for the Langevin equations[16,17,18,19].

Contrary to FT for classical systems, FT for quantum systems has not been well understood. In order to generalize FT to quantum systems, it is necessary to identify entropy change or work done externally. And two procedures are available:

- (a) Measure energy, particle numbers etc. twice and evaluate the entropy change or work done as the difference of the two observed values.
- (b) Measure flows of energy, particle numbers etc. and evaluate the entropy change or work done as accumulated values of the flows.

Needless to say, in classical systems, the two procedures give the same values of the entropy change or work done, because of the conservation of energy and particle numbers. However, in quantum systems, as a result of the noncommutativity of dynamical variables, the two procedures are not equivalent.

As far as the authors know, the procedure (a) was first suggested by Kurchan[20] and he showed that the fluctuation theorem (1) holds for the probability distribution function of the so-obtained entropy change or work done. Similar results were discussed by Hal Tasaki[21], Callens et al.[22], and the C^* generalization was given by Tasaki and Matsui[23].

On the other hand, the procedure (b) has not been well studied. In this paper, we investigate a quantum analogue of the fluctuation theorem, following the procedure (b), with respect to the work done externally for a harmonic oscillator linearly coupled with a harmonic reservoir. The oscillator is externally driven in such a way that the center of the harmonic potential follows a given trajectory. Note that the system is a quantum analogue of the Langevin equation[15,16,17,18,19] which describes the experiment by Wang et al.[14]. Then, we have found that (i) there appears a quantum correction of order $\hbar^2$ in TFT, and that (ii) the existence of a quantum correction for SSFT depends on the way of driving (no correction in the uniformly dragged case and $\hbar^2$ -correction in the periodically driven case). Moreover, the quantum correction for SSFT is universal in the sense that it depends only on the temperature and the frequency of the driving, but not on the specific features of the system and environment.

The rest of this paper is arranged as follows. The model is described in Sec.2. The equation of motion is solved and quantum analogues to the fluctuation theorems are derived in Sec.3. The subsequent sections are devoted to the discussions of TFT and SSFT separately. In the last section, discussions are given.

2 MODEL

In order to discuss quantum corrections to the fluctuation theorem, we study an exactly solvable model of a one-dimensional harmonic oscillator linearly coupled with a harmonic bath[24,25]. Because a straightforward derivation of a quantum Langevin equation is known[25], the Gardiner's version is used. Then, we consider the following thought experiment:

- (i) The system is prepared to be in equilibrium with inverse temperature β ¹.
- (ii) At time $t = 0$, the harmonic potential starts to move, where its center follows the trajectory $x = f(t)$.

When $t < 0$, the potential center is assumed to be fixed at $x = 0$ and to move continuously (namely, $f(0) = 0$). The Hamiltonian which describes the time

¹ Throughout this article, we use the unit where the Boltzmann constant is unity.

evolution is given as

$$H(t) = H_0 + \frac{k}{2}f(t)(-2q + f(t)) , \quad (2)$$

$$H_0 = \frac{p^2}{2m} + \frac{k}{2}q^2 + \frac{1}{2} : \int d\lambda ((p_\lambda - \kappa_\lambda q)^2 + \omega_\lambda^2 q_\lambda^2) : , \quad (3)$$

where q , p , and m stand for the coordinate, momentum and mass of the harmonic oscillator, respectively, k is the strength of the harmonic potential, q_λ and p_λ are the coordinate and momentum of a bath degree of freedom with frequency ω_λ , and $: \cdots :$ is the normal product with respect to the normal modes:

$$a_\lambda \equiv \frac{1}{\sqrt{2\hbar\omega_\lambda}}(\omega_\lambda q_\lambda + ip_\lambda) . \quad (4)$$

Here we assume that ω_λ runs from 0 to $+\infty$ and that the dispersion equation

$$\eta(z) \equiv mz^2 - k - \int d\lambda \kappa_\lambda^2 - \int d\lambda \frac{\omega_\lambda v_\lambda^2}{z^2 - \omega_\lambda^2} \quad (5)$$

has no real zeros, where

$$v_\lambda \equiv \sqrt{\omega_\lambda} \kappa_\lambda . \quad (6)$$

This condition corresponds to the case of the damped harmonic oscillator.

As in the previous works[14,16,17,15], we investigate the statistical property of the work externally done during the time interval τ divided by temperature β^{-1} : $\hat{\Sigma}_\tau$. If the work is started to be measured at time t , it is given by

$$\hat{\Sigma}_\tau \equiv \beta \int_0^\tau \dot{f}(s+t)(-k(q(s+t) - f(s+t)))ds \quad (7)$$

where $\dot{f}$ stands for the time derivative of f and $q(s+t)$ is the coordinate operator in the Heisenberg picture at time $s+t$. Then, the probability density $\pi_t(\Sigma_\tau = A)$ of the work is given by

$$\pi_t(\Sigma_\tau = A) = \langle e^{-\beta H_0} \delta(\hat{\Sigma}_\tau - A) \rangle \quad (8)$$

where $\langle \cdots \rangle \equiv \text{tr}(e^{-\beta H_0} \cdots)/Z$ stands for the thermal average and $Z = \text{tr} e^{-\beta H_0}$ is the partition function. Note that TFT refers to the symmetry of $\pi_0(\Sigma_\tau = A)$ while SSFT refers to that of $\pi_{+\infty}(\Sigma_\tau = A)$.

3 DISTRIBUTION FUNCTION OF THE WORK

In this section, we derive an explicit expression of the distribution function π_t of the work Σ_τ .

3.1 Evaluation of $\hat{\Sigma}_\tau$

First we note that the unperturbed Hamiltonian H_0 is diagonalized by the normal modes:

$$\alpha_\lambda = a_\lambda + \frac{v_\lambda}{\eta_+(\omega_\lambda)} \left\{ \frac{p - im\omega_\lambda q}{\sqrt{2\hbar}} + \int \frac{d\lambda'}{2} \left(\frac{v_{\lambda'} a_{\lambda'}}{\omega_\lambda - \omega_{\lambda'} + i0} - \frac{v_{\lambda'} a_{\lambda'}^+}{\omega_\lambda + \omega_{\lambda'}} \right) \right\} \quad (9)$$

where the normal mode a_λ and the interaction strength v_λ are given, respectively, by (4) and (6), $\eta_+(\omega_\lambda) \equiv \eta(\omega_\lambda + i0)$ and the function $\eta(z)$ is defined in (5). Indeed, one has $[\alpha_\lambda, \alpha_{\lambda'}^+] = \delta(\lambda - \lambda')$ and

$$H_0 = \int d\lambda \hbar \omega_\lambda \alpha_\lambda^+ \alpha_\lambda + (\text{c-number}) . \quad (10)$$

The old variables can be represented by these normal modes. For example,

$$q = \sqrt{\frac{\hbar}{2}} \int d\lambda \left(\frac{iv_\lambda}{\eta_-(\omega_\lambda)} \alpha_\lambda - \frac{iv_\lambda}{\eta_+(\omega_\lambda)} \alpha_\lambda^+ \right) \quad (11)$$

and, thus,

$$H(t) = H_0 - \sqrt{\frac{\hbar}{2}} k f(t) \int d\lambda \left\{ \frac{iv_\lambda \alpha_\lambda}{\eta_-(\omega_\lambda)} - \frac{iv_\lambda \alpha_\lambda^+}{\eta_+(\omega_\lambda)} \right\} + (\text{c-number}) \quad (12)$$

where $\eta_-(\omega_\lambda) = \eta(\omega_\lambda - i0)$.

From (12), one finds that the normal mode $\alpha_\lambda(t)$ in the Heisenberg picture obeys

$$\dot{\alpha}_\lambda(t) = -i\omega_\lambda \alpha_\lambda(t) + \frac{k f(t)}{\sqrt{2\hbar}} \frac{v_\lambda}{\eta_+(\omega_\lambda)} \quad (13)$$

which admits the solution:

$$\alpha_\lambda(t) = \alpha_\lambda e^{-i\omega_\lambda t} - i \frac{k v_\lambda}{\sqrt{2\hbar} \omega_\lambda \eta_+(\omega_\lambda)} \left(f(t) - \int_0^t e^{i\omega_\lambda(s-t)} \dot{f}(s) ds \right) \quad (14)$$

And we obtain the coordinate $q(t)$ in the Heisenberg picture as

$$q(t) = i \sqrt{\frac{\hbar}{2}} \int d\lambda \left(\frac{v_\lambda e^{-i\omega_\lambda t}}{\eta_-(\omega_\lambda)} \alpha_\lambda - \frac{v_\lambda e^{i\omega_\lambda t}}{\eta_+(\omega_\lambda)} \alpha_\lambda^+ \right) + k \int d\lambda \frac{v_\lambda^2}{|\eta_+(\omega_\lambda)|^2} \int_0^t \sin(\omega_\lambda(s-t)) f(s) ds \quad (15)$$

Then, by a tedious but straightforward calculation, one obtains the work operator $\hat{\Sigma}_\tau$

$$\hat{\Sigma}_\tau = -ik\beta \sqrt{\frac{\hbar}{2}} \left\{ \int d\lambda \frac{v_\lambda \hat{f}(\omega_\lambda; t, \tau)}{\eta_-(\omega_\lambda)} \alpha_\lambda - \int d\lambda \frac{v_\lambda \hat{f}^*(\omega_\lambda; t, \tau)}{\eta_+(\omega_\lambda)} \alpha_\lambda^+ \right\} + \int d\lambda \frac{\beta k^2 v_\lambda^2}{\omega_\lambda |\eta_+(\omega_\lambda)|^2} \int_t^{t+\tau} ds \int_0^s ds' \cos(\omega_\lambda(s-s')) \dot{f}(s) \dot{f}(s') \quad (16)$$

where $\hat{f}(\omega_\lambda; t, \tau)$ is a ‘partial Fourier transformation’ of $\dot{f}(t)$:

$$\hat{f}(\omega_\lambda; t, \tau) \equiv \int_t^{t+\tau} \dot{f}(s) e^{-i\omega_\lambda s} ds . \quad (17)$$

3.2 Calculation of the work distribution π_t

The distribution function $\pi_t(\Sigma_\tau = A)$ of the work $\hat{\Sigma}_\tau$ is obtained with the aid of the characteristic function

$$\Phi_t(\xi) = \int dA \pi_t(\Sigma_\tau = A) e^{i\xi A} = \langle \exp(i\xi \hat{\Sigma}_\tau) \rangle . \quad (18)$$

With the aid of the formula

$$\langle e^{\int d\lambda (\xi_\lambda \alpha_\lambda + \eta_\lambda \alpha_\lambda^+)} \rangle = e^{\int d\lambda \frac{\xi_\lambda \eta_\lambda}{2} \coth(\frac{\beta \hbar \omega_\lambda}{2})} \quad (19)$$

(its proof is given in Appendix), one obtains the following expression:

$$\Phi_t(\xi) = e^{i\xi m_{t,\tau} - \frac{\xi^2}{2} \sigma_{t,\tau}^2} , \quad (20)$$

where

$$m_{t,\tau} = \beta k^2 \int_t^{t+\tau} ds \int_0^s ds' \int d\lambda \frac{v_\lambda^2}{|\eta_+(\omega_\lambda)|^2} \frac{\cos(\omega_\lambda(s-s'))}{\omega_\lambda} \dot{f}(s) \dot{f}(s') \quad (21)$$

and

$$\sigma_{t,\tau}^2 = \frac{\hbar k^2 \beta^2}{2} \int_t^{t+\tau} ds \int_t^{t+\tau} ds' \int d\lambda \frac{v_\lambda^2 \cos(\omega_\lambda(s-s'))}{|\eta_+(\omega_\lambda)|^2} \coth\left(\frac{\beta \hbar \omega_\lambda}{2}\right) \dot{f}(s) \dot{f}(s') . \quad (22)$$

Eq.(20) shows that the distribution π_t is Gaussian with the mean value $m_{t,\tau}$ and the standard deviation $\sigma_{t,\tau}$:

$$\pi_t(\Sigma_\tau = A) = \frac{1}{\sqrt{2\pi}\sigma_{t,\tau}} \exp\left(-\frac{(A - m_{t,\tau})^2}{2\sigma_{t,\tau}^2}\right) . \quad (23)$$

By setting $t = 0$, (23) leads to the quantum counterpart of TFT[4,5]

$$\log \frac{\pi_0(\Sigma_\tau = A)}{\pi_0(\Sigma_\tau = -A)} = \frac{2m_{0,\tau}}{\sigma_{0,\tau}^2} A . \quad (24)$$

In addition, provided that the limits $m_{+\infty,\tau} = \lim_{t \rightarrow +\infty} m_{t,\tau}$ and $\sigma_{+\infty,\tau} = \lim_{t \rightarrow +\infty} \sigma_{t,\tau}$ exist, one has

$$\pi_{+\infty}(\Sigma_\tau = A) = \frac{1}{\sqrt{2\pi}\sigma_{+\infty,\tau}} \exp\left(-\frac{(A - m_{+\infty,\tau})^2}{2\sigma_{+\infty,\tau}^2}\right) . \quad (25)$$

where $\pi_{+\infty}(\Sigma_\tau = A) = \lim_{t \rightarrow +\infty} \pi_t(\Sigma_\tau = A)$, and the quantum counterpart of SSFT is derived

$$\lim_{\tau \rightarrow +\infty} \frac{1}{\tau} \log \frac{\pi_{+\infty}(\Sigma_\tau/\tau = a)}{\pi_{+\infty}(\Sigma_\tau/\tau = -a)} = \lim_{\tau \rightarrow +\infty} \frac{2m_{+\infty,\tau}}{\sigma_{+\infty,\tau}} a . \quad (26)$$

If the coefficient of A in (24) and that of a in (26) were unity, the fluctuation theorems would hold. However, the expressions of the mean value $m_{t,\tau}$ and the standard deviation $\sigma_{t,\tau}$ seem to indicate the deviations. In the following, we investigate this question for TFT and SSFT separately. In the latter case, since the existence of the $t \rightarrow +\infty$ limits strongly depends on the behavior of $f(t)$, we focus on two concrete examples: the uniformly dragged case $f(t) = v_0 t$ and the periodically driven case $f(t) = v_0/\Omega \sin \Omega t$.

4 QUANTUM CORRECTION FOR TFT

In this section, quantum corrections for TFT (transient fluctuation theorem) will be investigated.

First we note that $\sigma_{0,\tau}^2$ can be rewritten as

$$\sigma_{0,\tau}^2 = \hbar k^2 \beta^2 \int_0^\tau ds \int_0^s ds' \int d\lambda \frac{v_\lambda^2 \cos(\omega_\lambda(s-s'))}{|\eta_+(\omega_\lambda)|^2} \coth\left(\frac{\beta \hbar \omega_\lambda}{2}\right) \dot{f}(s) \dot{f}(s') . \quad (27)$$

As the integrand is different from that of $2m_{0,\tau}$, one has $\sigma_{0,\tau}^2 \neq 2m_{0,\tau}$ and, thus, TFT does not hold.

Using the expansion $\coth\left(\frac{\beta \hbar \omega_\lambda}{2}\right) = \frac{2}{\beta \hbar \omega_\lambda} + \frac{1}{6} \beta \hbar \omega_\lambda + O(\hbar^3)$, the quantum correction can be easily calculated. Indeed, one has

$$\begin{aligned} \sigma_{0,\tau}^2 &= \hbar k^2 \beta^2 \int_0^\tau ds \int_0^s ds' \int d\lambda \frac{v_\lambda^2}{|\eta_+(\omega_\lambda)|^2} \frac{\cos(\omega_\lambda(s-s'))}{\omega_\lambda} \dot{f}(s) \dot{f}(s') \\ &\quad + \frac{1}{12} \beta^3 \hbar^2 k^2 \int_0^\tau ds \int_0^s ds' \int d\lambda \frac{\omega_\lambda v_\lambda^2}{|\eta_+(\omega_\lambda)|^2} \cos(\omega_\lambda(s-s')) \dot{f}(s) \dot{f}(s') \\ &\quad + O(\hbar^4) \end{aligned} \quad (28)$$

and, thus,

$$\log \frac{\pi_0(\Sigma_\tau = A)}{\pi_0(\Sigma_\tau = -A)} = \left(1 + \hbar^2 \epsilon_2 + O(\hbar^4)\right) A . \quad (29)$$

where

$$\epsilon_2 = -\frac{\beta^3 k^2}{24 m_{0,\tau}} \int_0^\tau ds \int_0^s ds' \int d\lambda \frac{\omega_\lambda v_\lambda^2}{|\eta_+(\omega_\lambda)|^2} \cos(\omega_\lambda(s-s')) \dot{f}(s) \dot{f}(s') . \quad (30)$$

To illustrate the behavior of ϵ_2 , we consider the uniformly dragged case $f(t) = v_0 t$ where the damping is very weak: $\mu \equiv \int \kappa_\lambda^2 d\lambda / k \ll 1$. Up to the first order in μ , the function $1/\eta_\pm(\omega)$ can be approximated by

$$\frac{1}{\eta_\pm(\omega)} \simeq \frac{1}{m(\omega - \tilde{\Omega}_0 \pm i\gamma)(\omega + \tilde{\Omega}_0 \pm i\gamma)} \quad (31)$$

$$\tilde{\Omega}_0 = \sqrt{\frac{k}{m}} + \int \frac{d\lambda}{4m} \kappa_\lambda^2 \left\{ \text{P} \frac{1}{\sqrt{k/m} - \omega_\lambda} + \frac{1}{\sqrt{k/m} + \omega_\lambda} \right\} \quad (32)$$

$$\gamma = \frac{\pi}{4m} \int d\lambda \kappa_\lambda^2 \delta(\sqrt{k/m} - \omega_\lambda) \quad (33)$$

where the symbol P in the integrand stands for Cauchy's principal part. Then, by a straightforward calculation, one finds that the leading term of the correction ϵ_2 with respect to μ is independent of time τ and

$$\epsilon_2 = -\frac{\beta^2 k}{24m} (1 + \text{O}(\mu)) . \quad (34)$$

5 QUANTUM CORRECTION FOR SSFT

In this section, quantum corrections for SSFT (steady-state fluctuation theorem) will be investigated. In this case, the long-time limit $t \rightarrow +\infty$ should be taken. As the existence and the limiting value strongly depends on the potential motion $f(t)$, we examine quantum corrections for two examples: the uniformly dragged case $f(t) = v_0 t$ and the periodically driven case $f(t) = v_0/\Omega \sin \Omega t$.

5.1 Uniformly dragged case

In this case $f(t) = v_0 t$ and the mean work $m_{t,\tau}$ and its variance $\sigma_{t,\tau}^2$ read as

$$m_{t,\tau} = 2\beta k^2 v_0^2 \int d\lambda \frac{v_\lambda^2}{\omega_\lambda |\eta_+(\omega_\lambda)|^2} \frac{\sin(\omega_\lambda(t + \tau/2)) \sin \omega_\lambda \tau/2}{\omega_\lambda^2} \quad (35)$$

and

$$\sigma_{t,\tau}^2 = 2\hbar k^2 \beta^2 v_0^2 \int d\lambda \frac{v_\lambda^2}{|\eta_+(\omega_\lambda)|^2} \coth\left(\frac{\beta \hbar \omega_\lambda}{2}\right) \frac{\sin^2 \omega_\lambda \tau/2}{\omega_\lambda^2} . \quad (36)$$

In the limit of $t \rightarrow +\infty$, one finds

$$m_{+\infty,\tau} = \lim_{t \rightarrow +\infty} m_{t,\tau} = \int d\lambda \frac{2\pi \beta k^2 v_0^2 \kappa_\lambda^2}{|\eta_+(\omega_\lambda)|^2} \frac{\sin \omega_\lambda \tau/2}{\omega_\lambda} \delta(\omega_\lambda) = M_{+\infty} \tau \quad (37)$$

where $v_\lambda = \sqrt{\omega_\lambda} \kappa_\lambda$ is used and

$$M_{+\infty} = \frac{\pi \beta k^2 v_0^2}{|\eta_+(0)|^2} \int d\lambda \kappa_\lambda^2 \delta(\omega_\lambda) \quad (38)$$

Note that $\sigma_{+\infty, \tau}^2 = \sigma_{t, \tau}^2$ as the variance does not depend on t .

These equations together with $\{\sin^2 \omega \tau / 2\} / (\omega^2 \tau) \rightarrow \pi \delta(\omega) / 2$ ($\tau \rightarrow +\infty$) give

$$\begin{aligned} \lim_{\tau \rightarrow +\infty} \frac{\sigma_{+\infty, \tau}^2}{2m_{+\infty, \tau}} &= \frac{\pi \hbar k^2 \beta^2 v_0^2}{2M_{+\infty}} \int d\lambda \frac{\kappa_\lambda^2}{|\eta_+(\omega_\lambda)|^2} \omega_\lambda \coth\left(\frac{\beta \hbar \omega_\lambda}{2}\right) \delta(\omega_\lambda) \\ &= \frac{\pi \hbar k^2 \beta^2 v_0^2}{2M_{+\infty}} \frac{2}{\beta \hbar |\eta_+(0)|^2} \int d\lambda \kappa_\lambda^2 \delta(\omega_\lambda) = 1. \end{aligned} \quad (39)$$

Therefore, SSFT holds exactly:

$$\lim_{\tau \rightarrow +\infty} \frac{1}{\tau} \log \frac{\pi_{+\infty}(\Sigma_\tau / \tau = a)}{\pi_{+\infty}(\Sigma_\tau / \tau = -a)} = a. \quad (40)$$

The reason can be understood easily. Indeed, the motion of the potential is uniform in this case and only the zero frequency environmental modes do contribute to the steady-state work distribution. On the other hand, quantum effects appear only for the non-zero frequency modes and, thus, the quantum correction disappears in SSFT. In the next section, we consider a case where non-zero environmental modes do contribute to SSFT.

5.2 Periodically driven case

In this section, we consider the periodically driven case $\dot{f}(t) = v_0 \cos(\Omega t)$. First we remark that, in this case, there exists no exact steady state since the system is periodically driven. We consider, instead, a limiting oscillatory state obtained by a ‘stroboscopic limit’, where the time t is set to $t = 2\pi m / \Omega + \phi / \Omega$ and the limit $m \rightarrow +\infty$ is taken. The variable ϕ represents the phase of the limiting oscillation. Hereafter, we abbreviate

$$\lim_{m \rightarrow +\infty} F(t)|_{t=2\pi \frac{m}{\Omega} + \frac{\phi}{\Omega}} = \text{strob.} \lim_{t \rightarrow +\infty} F(t) \quad (41)$$

The variance of the work for $t = 2\pi m / \Omega + \phi / \Omega$ now reads as

$$\sigma_{t, \tau}^2 = \frac{\hbar k^2 \beta^2 v_0^2}{2} \int_0^\tau ds \int_0^\tau ds' \int d\lambda \frac{v_\lambda^2 \cos(\omega_\lambda(s - s'))}{|\eta_+(\omega_\lambda)|^2}$$

$$\times \coth\left(\frac{\beta\hbar\omega_\lambda}{2}\right) \cos(\Omega s + \phi) \cos(\Omega s' + \phi) . \quad (42)$$

Since it does not depend on m , one has $\sigma_{+\infty,\tau}^2 \equiv \text{strob.} \lim_{t \rightarrow +\infty} \sigma_{t,\tau}^2 = \sigma_{t,\tau}^2$. Then, by a tedious but straightforward calculation, one obtains

$$\lim_{\tau \rightarrow \infty} \sigma_{+\infty,\tau}^2 / \tau = \frac{\pi \hbar k^2 \beta^2 v_0^2}{4} \frac{\coth \frac{\beta \hbar \Omega}{2}}{|\eta_+(\Omega)|^2} \int d\lambda v_\lambda^2 \delta(\omega_\lambda - \Omega) . \quad (43)$$

As in the same way, one has

$$\begin{aligned} m_{+\infty,\tau} &= \text{strob.} \lim_{t \rightarrow +\infty} m_{t,\tau} \\ &= \beta k^2 v_0^2 \int_0^\tau ds \int_{-\infty}^s ds' \int d\lambda \frac{v_\lambda^2 \cos(\omega_\lambda(s-s'))}{\omega_\lambda |\eta_+(\omega_\lambda)|^2} \cos(\Omega s + \phi) \cos(\Omega s' + \phi) \\ &= \frac{\pi \beta k^2 v_0^2}{4} \frac{\tau}{\Omega |\eta_+(\Omega)|^2} \int d\lambda v_\lambda^2 \delta(\omega_\lambda - \Omega) \\ &\quad + \frac{\beta k^2 v_0^2}{4} \int d\lambda \frac{v_\lambda^2}{\omega_\lambda |\eta_+(\omega_\lambda)|^2} \left[P \frac{\cos 2\phi - \cos(2\Omega\tau + 2\phi)}{\Omega^2 - \omega_\lambda^2} \right. \\ &\quad \left. + \pi \delta(\Omega - \omega_\lambda) \frac{\sin(2\Omega\tau + 2\phi) - \sin 2\phi}{2\Omega} \right] \end{aligned} \quad (44)$$

and

$$\lim_{\tau \rightarrow +\infty} m_{+\infty,\tau} / \tau = \frac{\pi \beta k^2 v_0^2}{4\Omega |\eta_+(\Omega)|^2} \int d\lambda v_\lambda^2 \delta(\omega_\lambda - \Omega) . \quad (45)$$

Eqs.(43) and (45) give

$$\lim_{\tau \rightarrow +\infty} \frac{2m_{+\infty,\tau}}{\sigma_{+\infty,\tau}^2} = \frac{2}{\beta \hbar \Omega} \tanh \frac{\beta \hbar \Omega}{2} \quad (46)$$

and, thus, SSFT does not hold

$$\lim_{\tau \rightarrow +\infty} \frac{1}{\tau} \log \frac{\pi_{+\infty}(\Sigma_\tau / \tau = a)}{\pi_{+\infty}(\Sigma_\tau / \tau = -a)} = \frac{2}{\beta \hbar \Omega} \tanh \frac{\beta \hbar \Omega}{2} a \neq a . \quad (47)$$

where the distribution function refers to the limiting oscillatory state. Note that the quantum correction is universal in the sense that it depend only on the temperature and the frequency of the driving, but not on the specific features of the system and environment.

6 DISCUSSIONS

In summary, for a harmonic oscillator linearly coupled with a harmonic reservoir, we have investigated quantum analogues of the transient and steady-state fluctuation theorems with respect to the work done externally. And we have shown the followings:

- (i) In the transient fluctuation theorem, there appears a quantum correction of order $\hbar^2$.
- (ii) In the steady-state fluctuation theorem, the existence of a quantum correction depends on the way of driving. In the uniformly dragged case, the classical formula holds, while, in the periodically driven case, there appears a correction of order $\hbar^2$.

We remind that the steady-state fluctuation theorem for the periodically driven case refers to the limiting oscillatory state, which is obtained as the stroboscopic limit defined in (41).

Remarkably, the situation is in contrast to that for Kurchan’s quantum fluctuation theorem, where energy is measured twice, and the entropy change and work done are evaluated from the difference between the two measured values. In this case, as discussed previously[20,21,22,23], the (transient) fluctuation theorem holds as in the classical systems.

Finally, we note that the quantum correction for the steady-state fluctuation theorem is universal in the sense that it depends only on the temperature and the frequency of the driving, but not on the specific features of the system and environment. We expect that this is also the case for general systems.

Acknowledgement

The authors are grateful to Professors T. Matsui and M. Sano for fruitful discussions and valuable comments. This work is supported by a Grant-in-Aid for Scientific Research (C) from JSPS, by a Grant-in-Aid for Scientific Research of Priority Areas “Control of Molecules in Intense Laser Fields” and 21st Century COE Program (Holistic research and Education Center for Physics of Self-Organization Systems) both from the Ministry of Education, Culture, Sports, Science and Technology of Japan.

A Derivation of $\langle e^{\int d\lambda(\xi_\lambda \alpha_\lambda + \eta_\lambda \alpha_\lambda^+)} \rangle = e^{\int d\lambda \frac{\xi_\lambda \eta_\lambda}{2} \coth(\frac{\beta \hbar \omega}{2})}$.

The desired result immediately follows from

$$\langle e^{\eta a^+ + \xi a} \rangle = e^{\frac{\eta \xi}{2} \coth(\frac{\beta \hbar \omega}{2})}, \quad (\text{A.1})$$

where $[a, a^+] = 1$ and $\langle \dots \rangle \equiv \text{tr}(\dots e^{-\beta \hbar \omega a^+ a}) / Z$ with $Z = \text{tr}(e^{-\beta \hbar \omega a^+ a})$. This relation can be shown as follows.

The Baker-Hausdorff theorem gives

$$e^{\eta a^+ + \xi a} = e^{\eta a^+} e^{\xi a} e^{\frac{\xi \eta}{2}}. \quad (\text{A.2})$$

With the aid of the coherent states $|\alpha\rangle \equiv \sum_{n=0}^{\infty} \frac{\alpha^n}{\sqrt{n!}} |n\rangle$, which satisfy

$$a |\alpha\rangle = \alpha |\alpha\rangle, \quad \langle \alpha | a^+ = \alpha^* \langle \alpha | \quad (\text{A.3})$$

$$\int \frac{d\alpha}{\pi} |\alpha\rangle \langle \alpha| e^{-|\alpha|^2} = 1, \quad \langle \alpha | \beta \rangle = e^{\alpha^* \beta}, \quad (\text{A.4})$$

one finds

$$\begin{aligned} \langle e^{\eta a^+ + \xi a} \rangle &= e^{\frac{\xi \eta}{2}} \langle e^{\eta a^+} e^{\xi a} \rangle = \frac{e^{\frac{\xi \eta}{2}}}{Z} \text{tr}(e^{-\beta \hbar \omega a^+ a} e^{\eta a^+} e^{\xi a}) \\ &= \frac{e^{\frac{\xi \eta}{2}}}{Z} \sum_{n=0}^{\infty} e^{-\beta \hbar \omega n} \langle n | \int \frac{d\alpha}{\pi} |\alpha\rangle \langle \alpha| e^{\eta a^+} e^{\xi a} \int \frac{d\beta'}{\pi} |\beta'\rangle \langle \beta'| n \rangle e^{-|\alpha|^2 - |\beta'|^2} \\ &= \frac{e^{\frac{\xi \eta}{2}}}{Z} \sum_{n=0}^{\infty} e^{-\beta \hbar \omega n} \int \frac{d\alpha}{\pi} \int \frac{d\beta'}{\pi} e^{-|\alpha|^2 - |\beta'|^2} \langle \alpha | \beta' \rangle e^{\eta \alpha^*} e^{\xi \beta'} \frac{\alpha^n \beta'^{*n}}{n!} \\ &= \frac{e^{\frac{\xi \eta}{2}}}{Z} \int \frac{d\alpha}{\pi} \int \frac{d\beta'}{\pi} e^{-|\alpha|^2 - |\beta'|^2 + \xi \beta' + \eta \alpha^* + \alpha \beta'^*} e^{-\beta \hbar \omega + \alpha^* \beta'} \\ &= e^{\frac{\xi \eta}{2}} \frac{1}{Z} \int \frac{d\beta'}{\pi} e^{-(1 - e^{-\beta \hbar \omega}) |\beta'|^2 + e^{-\beta \hbar \omega} \beta'^* \eta + \beta' \xi} \\ &= e^{\frac{\eta \xi}{2} \coth(\frac{\beta \hbar \omega}{2})}, \end{aligned} \quad (\text{A.5})$$

which leads to the desired relation (A.1):

$$\langle e^{\eta a^+ + \xi a} \rangle = e^{\frac{\eta \xi}{2} \coth(\frac{\beta \hbar \omega}{2})} \quad (\text{A.6})$$




Fig. B.1. Integration contour for the integral (B.3)

B Integral containing $\eta_+(\omega_\lambda)$

To proceed the calculations in Sec.3, one has to evaluate integrals involving $\eta_+(\omega_\lambda)$. In this appendix, we illustrate the way by evaluating

$$\int d\lambda \frac{v_\lambda^2}{\omega_\lambda |\eta_+(\omega_\lambda)|^2} .$$

The key idea is to rewrite it as a contour integral and to use the relations:

$$\eta_-(x) - \eta_+(x) = -\pi i \int d\lambda \delta(x - \omega_\lambda) v_\lambda^2 , \quad (\text{B.1})$$

$$\eta_\pm(-x) = \eta_\mp(x) . \quad (\text{B.2})$$

Then, one has

$$\begin{aligned} \int d\lambda \frac{v_\lambda^2}{\omega_\lambda |\eta_+(\omega_\lambda)|^2} &= \int d\lambda \int_0^\infty dx \delta(x - \omega_\lambda) \frac{v_\lambda^2}{\eta_-(x)\eta_+(x)} \frac{1}{x} \\ &= \int_0^\infty \frac{1}{-\pi i} \left(\frac{1}{\eta_+(x)} - \frac{1}{\eta_-(x)} \right) \frac{1}{x} dx = \int_{-\infty}^\infty \frac{1}{-\pi i} \frac{1}{\eta_+(x)} \frac{1}{x} dz \\ &= \int_C \frac{1}{-\pi i} \frac{dz}{z\eta(z)} - \lim_{r \rightarrow 0} \int_0^{2\pi} d\theta \frac{1}{-\pi i} \frac{r i e^{i\theta}}{r e^{i\theta} \eta(r e^{i\theta})} - \lim_{R \rightarrow \infty} \int_0^{2\pi} \frac{1}{-\pi i} \frac{R i e^{i\theta} d\theta}{R e^{i\theta} \eta(R e^{i\theta})} \\ &= \frac{1}{k} , \end{aligned} \quad (\text{B.3})$$

where the integration contour C is shown in Fig.B.1.

References

- [1] D.J. Evans, E.G.D. Cohen and G.P. Morriss, Phys. Rev. Lett. 71 (1993) 2401.
- [2] C. Jarzynski, Phys. Rev. Lett. 78 (1996) 2690; J. Stat. Phys. 95 (1999) 367.
- [3] G. Gallavotti and E.G.D. Cohen, Phys. Rev. Lett. 74 (1995) 2694; J. Stat. Phys. 80 (1995) 931; E.G.D. Cohen and G. Gallavotti, J.Stat.Phys. 96 (1999) 1343.
- [4] D.J. Evans and D. Searles, Phys. Rev. E 50 (1994) 1645.
- [5] D. Searles and D.J. Evans, J. Chem. Phys. 112 (2000) 9727; *ibid.* 113 (2000) 3503; D.J. Evans and D. Searles, Adv. Phys. 51 (2002) 1529; E. Mittag and D.J. Evans, Phys. Rev. E 67 (2003) 026113.
- [6] J. Kurchan, J. Phys. A 31 (1998) 3719.
- [7] J.L. Lebowitz and H. Spohn, J. Stat. Phys. 95 (1999) 333.
- [8] C. Maes, J. Stat. Phys. 95 (1999) 367.
- [9] C. Jarzynski, J. Stat. Phys. 98 (2000) 77.
- [10] L. Rondoni, T. Tél and J. Vollmer, Phys. Rev. E 61 (2000) R4679.
- [11] L. Rey-Bellet and L.E. Thomas, Ann. H. Poincaré 3 (2002) 483.
- [12] G.E. Crooks, Phys. Rev. E 60 (1999) 2721.
- [13] C. Maes and K. Netočný, J. Stat. Phys. 110 (2003) 269.
- [14] G.M. Wang, E.M. Sevick, E. Mittag, D.J. Evans and D. Searles, Phys. Rev. Lett. 89 (2002) 050601.
- [15] O. Mazonka and C. Jarzynski, e-print cond-mat/9912121.
- [16] S.Tasaki, I.Terasaki and T.Monnai, e-print cond-mat/0208153v2.
- [17] R. van Zon and E.G.D. Cohen, Phys. Rev. E 67 (2003) 046102.
- [18] R. van Zon and E.G.D. Cohen, e-print cond-mat/0305147v1; e-print cond-mat/0307297v1.
- [19] O. Narayan and A. Dhar, e-print cond-mat/0307148v1.
- [20] J. Kurchan, e-print cond-mat/0007360v2.
- [21] Hal Tasaki, e-print cond-mat/0009244.
- [22] I. Callens, W. De Roeck, T. Jacobs, C. Maes and K. Netočný, e-print cond-mat/0211252v2.
- [23] S. Tasaki and T. Matsui, in: Fundamental Aspects of Quantum Physics, eds. L. Accardi and S. Tasaki (World Scientific, Singapore, 2003) p.100.
- [24] G.W. Ford, M. Kac and P. Mazur, J. Math. Phys. 6 (1965) 504.
- [25] C.W.Gardiner, Quantum Noise, (Springer Verlag, Berlin, 1991) Chap.3.